

2005 AIME I Solutions

Typeset by: LIVE by Po-Shen Loh

<https://live.poshenloh.com/past-contests/aime/2005I/solutions>



Problems © Mathematical Association of America. Reproduced with permission.

1. Six congruent circles form a ring with each circle externally tangent to the two circles adjacent to it. All six circles are internally tangent to a circle $\mathcal{C}$ with radius 30. Let K be the area of the region inside $\mathcal{C}$ and outside all of the six circles in the ring. Find $\lfloor K \rfloor$. (The notation $\lfloor K \rfloor$ denotes the greatest integer that is less than or equal to K .)



Solution:

Let r be the common radius of the six circles. Adjacent circles are externally tangent, so their centers are $2r$ apart, and the six centers form a regular hexagon with side $2r$. Since a regular hexagon's circumradius equals its side length, each center is at distance $2r$ from the center O of $\mathcal{C}$. Internal tangency to $\mathcal{C}$ means the distance from O to each small center plus r equals 30, so $3r = 30$ and $r = 10$.

Therefore

$$\begin{aligned} K &= \pi (30^2 - 6 \cdot 10^2) \\ &= 300\pi \approx 942.48, \end{aligned}$$

and $\lfloor K \rfloor = 942$.

2. For each positive integer k , let S_k denote the increasing arithmetic sequence of integers whose first term is 1 and whose common difference is k . For example, S_3 is the sequence $1, 4, 7, \dots$. For how many values of k does S_k contain the term 2005?



Solution:

The n th term of S_k is $1 + (n - 1)k$, so 2005 is a term exactly when $(n - 1)k = 2004$ for some positive integer n , that is, exactly when k divides 2004. Every divisor works, since $n - 1 = \frac{2004}{k}$ is then a positive integer.

Since $2004 = 2^2 \cdot 3 \cdot 167$, the number of divisors is $(2 + 1)(1 + 1)(1 + 1) = 12$.

3. How many positive integers have exactly three proper divisors, each of which is less than 50? (A *proper divisor* of a positive integer n is a positive integer divisor of n other than n itself.)



Solution:

An integer with exactly three proper divisors has exactly four divisors in total, so it is either $n = pq$ with p and q distinct primes (proper divisors $1, p, q$) or $n = p^3$ with p prime (proper divisors $1, p, p^2$).

In the first case we need p and q both less than 50. There are 15 primes below 50, giving $\binom{15}{2} = 105$ such numbers. In the second case we need $p^2 < 50$, which holds for $p = 2, 3, 5, 7$, giving 4 more.

The total is $105 + 4 = 109$.

4. The director of a marching band wishes to place the members into a formation that includes all of them and has no unfilled positions. If they are arranged in a square formation, there are 5 members left over. The director finds that if they are arranged in a rectangular formation with 7 more rows than columns, the desired result can be obtained. Find the maximum number of members this band can have.



Solution:

Let the band have n members, with $n = s^2 + 5$ for the square formation and $n = x(x + 7)$ for the rectangular formation with x columns. Multiplying $x^2 + 7x = s^2 + 5$ by 4 and completing the square gives

$$(2x + 7)^2 - (2s)^2 = 69,$$

so $(2x + 7 - 2s) \cdot (2x + 7 + 2s) = 69$.

Writing $69 = 1 \cdot 69 = 3 \cdot 23$ with the larger factor second: from $1 \cdot 69$ we get $2x + 7 = 35$ and $2s = 34$, so $x = 14$, $s = 17$, and $n = 17^2 + 5 = 294$. From $3 \cdot 23$ we get $2x + 7 = 13$ and $2s = 10$, so $x = 3$, $s = 5$, and $n = 30$.

The maximum is 294, achieved by a 21×14 rectangle.

5. Robert has 4 indistinguishable gold coins and 4 indistinguishable silver coins. Each coin has an engraving of a face on one side, but not on the other. He wants to stack the eight coins on a table into a single stack so that no two adjacent coins are face to face. Find the number of possible distinguishable arrangements of the 8 coins.



Solution:

Choose the coin orientations and the gold/silver positions independently. Record the orientations from bottom to top as a string of U (engraved face up) and D (engraved face down). Two adjacent coins are face to face exactly when the lower coin's engraved side faces up and the upper coin's engraved side faces down — that is, exactly when a U is immediately followed by a D.

A string of U's and D's avoids the pattern UD exactly when every D precedes every U, so the string is $D^i U^{8-i}$ for some $i = 0, 1, \dots, 8$: there are 9 allowable orientation strings. Independently, the gold coins occupy 4 of the 8 positions in $\binom{8}{4} = 70$ ways.

The total is $9 \cdot 70 = 630$.

6. Let P be the product of the nonreal roots of $x^4 - 4x^3 + 6x^2 - 4x = 2005$. Find $\lfloor P \rfloor$.
(The notation $\lfloor P \rfloor$ denotes the greatest integer that is less than or equal to P .)



Solution:

Adding 1 to both sides turns the left side into a perfect fourth power:

$$\begin{aligned}(x - 1)^4 &= x^4 - 4x^3 + 6x^2 \\ &\quad - 4x + 1 = 2006.\end{aligned}$$

So $x - 1$ is a fourth root of 2006 : the four roots are $x = 1 \pm \sqrt[4]{2006}$ (real) and $x = 1 \pm i\sqrt[4]{2006}$ (nonreal).

The product of the conjugate pair of nonreal roots is

$$\begin{aligned}&\left(1 + i\sqrt[4]{2006}\right) \\ &\quad \cdot \left(1 - i\sqrt[4]{2006}\right) \\ &= 1 + \sqrt{2006}.\end{aligned}$$

Since $44^2 = 1936 < 2006 < 2025 = 45^2$, we have $45 < P < 46$, so $\lfloor P \rfloor = 45$.

7. In quadrilateral $ABCD$, $BC = 8$, $CD = 12$, $AD = 10$, and $m\angle A = m\angle B = 60^\circ$. Given that $AB = p + \sqrt{q}$, where p and q are positive integers, find $p + q$.



Solution:

Extend rays AD and BC until they meet at P . Triangle ABP has 60° angles at A and B , so it is equilateral: $PA = PB = AB$. Writing $x = AB$, we get $PD = PA - AD = x - 10$ and $PC = PB - BC = x - 8$.

The Law of Cosines in triangle PDC , with $\angle P = 60^\circ$ and $DC = 12$, gives

$$\begin{aligned} 144 &= (x - 10)^2 + (x - 8)^2 \\ &\quad - (x - 10)(x - 8) \\ &= x^2 - 18x + 84, \end{aligned}$$

so $x^2 - 18x - 60 = 0$ and $x = 9 + \sqrt{81 + 60} = 9 + \sqrt{141}$.

Thus $p + q = 9 + 141 = 150$.

8. The equation

$$2^{333x-2} + 2^{111x+2} = 2^{222x+1} + 1$$

has three real roots. Given that their sum is $\frac{m}{n}$, where m and n are relatively prime positive integers, find $m + n$.



Solution:

Let $y = 2^{111x}$. Then $2^{333x-2} = \frac{y^3}{4}$, $2^{111x+2} = 4y$, and $2^{222x+1} = 2y^2$, so the equation becomes $\frac{y^3}{4} + 4y = 2y^2 + 1$, that is,

$$y^3 - 8y^2 + 16y - 4 = 0.$$

Since the three roots x_1, x_2, x_3 are real and $y = 2^{111x}$ is strictly increasing, they correspond to three positive real roots y_1, y_2, y_3 of the cubic.

Each $x_i = \frac{1}{111} \log_2 y_i$, so

$$\begin{aligned} x_1 + x_2 + x_3 &= \frac{1}{111} \log_2(y_1 y_2 y_3) \\ &= \frac{1}{111} \log_2 4 \\ &= \frac{2}{111}, \end{aligned}$$

using Vieta's formulas for the product of the roots. Then $m + n = 2 + 111 = 113$.

9. Twenty-seven unit cubes are each painted orange on a set of four faces so that the two unpainted faces share an edge. The 27 cubes are then randomly arranged to form a $3 \times 3 \times 3$ cube. Given that the probability that the entire surface of the larger cube is orange is $\frac{p^a}{q^b r^c}$, where p, q , and r are distinct primes and a, b , and c are positive integers, find $a + b + c + p + q + r$.



Solution:

Each unit cube has one “bad edge”: the edge shared by its two unpainted faces. The larger cube’s surface is entirely orange exactly when every unit cube’s bad edge touches no visible face. A uniformly random orientation places the bad edge uniformly among the cube’s 12 edge positions, so for each unit cube we count the edge positions both of whose faces are hidden.

A corner cube shows 3 faces meeting at a vertex; the safe edges are those of the 3 hidden faces meeting at the opposite vertex, so the probability is $\frac{3}{12} = \frac{1}{4}$. An edge cube shows 2 adjacent faces, which touch $4 + 4 - 1 = 7$ edges, leaving 5 safe: probability $\frac{5}{12}$. A face-center cube shows 1 face touching 4 edges, leaving 8 safe: probability $\frac{8}{12} = \frac{2}{3}$. The center cube is always fine.

With 8 corner, 12 edge, and 6 face-center cubes, the probability is

$$\left(\frac{1}{4}\right)^8 \left(\frac{5}{12}\right)^{12} \left(\frac{2}{3}\right)^6$$

$$= \frac{5^{12}}{2^{34} \cdot 3^{18}},$$

so $a + b + c + p + q + r = 12 + 34 + 18 + 5 + 2 + 3 = 74$.

10. Triangle ABC lies in the Cartesian plane and has area 70. The coordinates of B and C are $(12, 19)$ and $(23, 20)$, respectively, and the coordinates of A are (p, q) . The line containing the median to side $\overline{BC}$ has slope -5 . Find the largest possible value of $p + q$.



Solution:

The median to $\overline{BC}$ passes through the midpoint $M = \left(\frac{35}{2}, \frac{39}{2}\right)$ of $\overline{BC}$. The line through M with slope -5 is $y = -5x + 107$, and A lies on this line, so $A = (p, -5p + 107)$ and $q = -5p + 107$.

By the shoelace formula with $B = (12, 19)$ and $C = (23, 20)$,

$$\begin{aligned}[ABC] &= \frac{1}{2} |-p + 12(20 - q) + 23(q - 19)| \\ &= \frac{1}{2} |980 - 56p| = 70,\end{aligned}$$

so $|56p - 980| = 140$, giving $p = 15$ or $p = 20$.

Since $p + q = p + (-5p + 107) = 107 - 4p$, the smaller value $p = 15$ gives the larger sum $107 - 60 = 47$.

11. A semicircle with diameter d is contained in a square whose sides have length 8. Given that the maximum value of d is $m - \sqrt{n}$, where m and n are integers, find $m + n$.



Solution:

Scale to a semicircle of radius 1 and ask for the smallest square containing it when its diameter makes angle θ with one pair of sides, where $0 \leq \theta \leq 90^\circ$. Squeeze the semicircle between two pairs of parallel lines in the square's two side directions: in each direction one line of the pair is tangent to the arc and the other passes through an endpoint of the diameter, and the distances between the pairs are $1 + \cos \theta$ and $1 + \sin \theta$. So the smallest enclosing square in that orientation has side $\max\{1 + \cos \theta, 1 + \sin \theta\}$, which is minimized when $\theta = 45^\circ$, giving side $1 + \frac{\sqrt{2}}{2} = \frac{2+\sqrt{2}}{2}$.

Scaling this optimal configuration so the square has side 8, the radius becomes $r = \frac{8}{\frac{2+\sqrt{2}}{2}} = \frac{16}{2+\sqrt{2}} = 8(2 - \sqrt{2})$, so

$$\begin{aligned} d = 2r &= 16(2 - \sqrt{2}) \\ &= 32 - 16\sqrt{2} = 32 - \sqrt{512}. \end{aligned}$$

Thus $m + n = 32 + 512 = 544$.

12. For positive integers n , let $\tau(n)$ denote the number of positive integer divisors of n , including 1 and n . For example, $\tau(1) = 1$ and $\tau(6) = 4$. Define $S(n)$ by

$$S(n) = \tau(1) + \tau(2) + \cdots + \tau(n).$$

Let a denote the number of positive integers $n \leq 2005$ with $S(n)$ odd, and let b denote the number of positive integers $n \leq 2005$ with $S(n)$ even. Find $|a - b|$.



Solution:

Divisors of n pair up as d and $\frac{n}{d}$, so $\tau(n)$ is odd exactly when n is a perfect square. Hence $S(n)$ changes parity exactly at the squares, which means $S(n)$ is odd exactly when the number of squares up to n , namely $\lfloor \sqrt{n} \rfloor$, is odd.

For each k , there are $2k + 1$ integers n with $\lfloor \sqrt{n} \rfloor = k$, namely $k^2 \leq n \leq k^2 + 2k$. Since $44^2 = 1936 \leq 2005 < 2025 = 45^2$, the odd values $k = 1, 3, \dots, 43$ all have their full blocks within range, so

$$\begin{aligned} a &= \sum_{k \text{ odd}, k \leq 43} (2k + 1) \\ &= 2(1 + 3 + \cdots + 43) + 22 \\ &= 2 \cdot 484 + 22 \\ &= 990. \end{aligned}$$

Then $b = 2005 - 990 = 1015$, and $|a - b| = 25$.

13. A particle moves in the Cartesian plane from one lattice point to another according to the following rules:

- From any lattice point (a, b) , the particle may move only to $(a + 1, b)$, $(a, b + 1)$, or $(a + 1, b + 1)$.
- There are no right angle turns in the particle's path. That is, the sequence of points visited contains neither a subsequence of the form $(a, b), (a + 1, b), (a + 1, b + 1)$ nor a subsequence of the form $(a, b), (a, b + 1), (a + 1, b + 1)$.

How many different paths can the particle take from $(0, 0)$ to $(5, 5)$?



Solution:

The forbidden right-angle turns say exactly that a rightward step may never immediately follow an upward step, and vice versa; a diagonal step may follow or precede anything. So at each lattice point (x, y) track three counts $D(x, y), R(x, y), U(x, y)$: the numbers of legal paths from $(0, 0)$ arriving there by a diagonal, rightward, or upward step. The rules give

$$\begin{aligned} D(x, y) &= D + R \\ &\quad + U \text{ at } (x - 1, y - 1), \\ R(x, y) &= D(x - 1, y) \\ &\quad + R(x - 1, y), \\ U(x, y) &= D(x, y - 1) \\ &\quad + U(x, y - 1). \end{aligned}$$

Starting from the single empty path at $(0, 0)$ (which may begin with any step), fill in the grid up to $(5, 5)$. Along the axes only all-rightward or all-upward paths survive, and the interior builds up quickly; at $(5, 5)$ the three counts come out to 27, 28, and 28.

The total number of paths is $27 + 28 + 28 = 83$.

14. Consider the points $A(0, 12)$, $B(10, 9)$, $C(8, 0)$, and $D(-4, 7)$. There is a unique square $\mathcal{S}$ such that each of the four points is on a different side of $\mathcal{S}$. Let K be the area of $\mathcal{S}$. Find the remainder when $10K$ is divided by 1000.



Solution:

Since segments $\overline{AC}$ and $\overline{BD}$ cross, A and C lie on opposite sides of the square, as do B and D . Let m be the slope of the side through B , so that side lies on $mx - y + 9 - 10m = 0$, and the perpendicular side through C lies on $x + my - 8 = 0$. The side length of the square equals both the distance between the parallel sides through B and D and the distance between the sides through A and C :

$$\begin{aligned} & \frac{|-4m - 7 + 9 - 10m|}{\sqrt{m^2 + 1}} \\ &= \frac{|12m - 8|}{\sqrt{m^2 + 1}}, \end{aligned}$$

so $|2 - 14m| = |12m - 8|$, giving $m = \frac{5}{13}$ or $m = -3$.

For $m = \frac{5}{13}$, the points A and C fall on opposite sides of the line through B , which is impossible if that line contains a side of the square, so $m = -3$. Then the side length is $\frac{|12(-3) - 8|}{\sqrt{9+1}} = \frac{44}{\sqrt{10}}$, so

$$K = \frac{44^2}{10} = \frac{1936}{10}$$

and

$$10K = 1936.$$

The remainder when 1936 is divided by 1000 is 936.

15. In $\triangle ABC$, $AB = 20$. The incircle of the triangle divides the median containing C into three segments of equal length. Given that the area of $\triangle ABC$ is $m\sqrt{n}$, where m and n are integers and n is not divisible by the square of any prime, find $m + n$.



Solution:

Let M be the midpoint of $\overline{AB}$, and let the incircle cut median $\overline{CM}$ at S and N , with $CS = SN = NM = \frac{1}{3}CM$. Let the incircle touch $\overline{AB}$ at T and $\overline{AC}$ at R . By Power of a Point,

$$\begin{aligned} MT^2 &= MN \cdot MS \\ &= \frac{CM}{3} \cdot \frac{2CM}{3} \\ &= \frac{2}{9}CM^2, \\ CR^2 &= CS \cdot CN \\ &= \frac{2}{9}CM^2, \end{aligned}$$

so $MT = CR$. Since $AR = AT$ (tangents from A), we get $AC = AR + RC = AT + TM = AM = 10$.

Write $a = BC$ and $s = \frac{20+a+10}{2} = 15 + \frac{a}{2}$. The standard tangent length gives $AT = s - a$, so $MT = AM - AT = 10 - (15 - \frac{a}{2}) = \frac{a-10}{2}$, while the median length formula gives $CM^2 = \frac{2 \cdot 10^2 + 2a^2 - 20^2}{4} = \frac{a^2 - 100}{2}$. Substituting into $MT^2 = \frac{2}{9}CM^2$:

$$\begin{aligned} \frac{(a-10)^2}{4} &= \frac{a^2 - 100}{9} \\ &\implies 9(a-10) \\ &= 4(a+10) \\ &\implies a = 26. \end{aligned}$$

Then the sides are 20, 26, 10 with $s = 28$, and Heron's formula gives

$$\begin{aligned}[ABC] &= \sqrt{28 \cdot 8 \cdot 2 \cdot 18} \\ &= \sqrt{8064} = 24\sqrt{14},\end{aligned}$$

so $m + n = 24 + 14 = 38$.

Problems: <https://live.poshenloh.com/past-contests/aime/2005I>

