

2025 AMC 12A Solutions

Typeset by: LIVE by Po-Shen Loh

<https://live.poshenloh.com/past-contests/amc12/2025A/solutions>



Problems © Mathematical Association of America. Reproduced with permission.

1. Andy and Betsy both live in Mathville. Andy leaves Mathville on his bicycle at 1:30, traveling due north at a steady 8 miles per hour. Betsy leaves on her bicycle from the same point at 2:30, traveling due east at a steady 12 miles per hour. At what time will they be exactly the same distance from their common starting point?

A 3:30

B 3:45

C 4:00

D 4:15

E 4:30

Solution:

Let t be the number of hours since 1:30. Andy has traveled $8t$ miles north, and Betsy, who started an hour later, has traveled $12(t - 1)$ miles east.

Setting the distances equal,

$$8t = 12(t - 1),$$

so $4t = 12$ and $t = 3$.

Three hours after 1:30 is 4:30.

Thus, the correct answer is **E**.

2. A box contains 10 pounds of a nut mix that is 50 percent peanuts, 20 percent cashews, and 30 percent almonds. A second nut mix containing 20 percent peanuts, 40 percent cashews, and 40 percent almonds is added to the box resulting in a new nut mix that is 40 percent peanuts. How many pounds of cashews are now in the box?

A 3.5

B 4

C 4.5

D 5

E 6

Solution:

The first box has 5 lb peanuts, 2 lb cashews, and 3 lb almonds. Adding x pounds of the second mix contributes $0.2x$ lb peanuts and $0.4x$ lb cashews.

The new peanut fraction is 40%, so

$$\frac{5 + 0.2x}{10 + x} = 0.4.$$

This gives $5 + 0.2x = 4 + 0.4x$, so $x = 5$.

The cashews now total $2 + 0.4(5) = 4$ pounds.

Thus, the correct answer is **B**.

3. A team of students is going to compete against a team of teachers in a trivia contest. The total number of students and teachers is 15. Ash, a cousin of one of the students, wants to join the contest. If Ash plays with the students, the average age on that team will increase from 12 to 14. If Ash plays with the teachers, the average age on that team will decrease from 55 to 52. How old is Ash?

A 28

B 29

C 30

D 32

E 33

Solution:

Let s be the number of students and a be Ash's age. The students' ages total $12s$, and adding Ash gives

$$\begin{aligned}12s + a &= 14(s + 1) \\ \implies a &= 2s + 14.\end{aligned}$$

There are $15 - s$ teachers with ages totaling $55(15 - s)$, and adding Ash gives

$$\begin{aligned}55(15 - s) + a &= 52(16 - s) \\ \implies a &= 3s + 7.\end{aligned}$$

Setting $2s + 14 = 3s + 7$ gives $s = 7$, so $a = 2(7) + 14 = 28$.

Thus, the correct answer is **A**.

4. Agnes writes the following four statements on a blank piece of paper.

- At least one of these statements is true.
- At least two of these statements are true.
- At least two of these statements are false.
- At least one of these statements is false.

Each statement is either true or false. How many false statements did Agnes write on the paper?

- ☐ A 0
- ☒ B 1
- ☐ C 2
- ☐ D 3
- ☐ E 4

Solution:


Let T be the number of true statements. The statements assert $T \geq 1, T \geq 2, T \leq 2$, and $T \leq 3$, respectively.

Testing $T = 3$: the conditions $T \geq 1, T \geq 2, T \leq 3$ hold (statements one, two, four) and $T \leq 2$ fails (statement three). Exactly 3 statements are true, matching $T = 3$.

For $T = 0, 1, 2, 3, 4$, the respective numbers of true conditions are 2, 3, 4, 3, 2, so no other value is self-consistent. Therefore exactly one statement is false.

Thus, the correct answer is **B**.

5. In the figure below, the outside square contains infinitely many squares, each of them with the same center and sides parallel to the outside square. The ratio of the side length of a square to the side length of the next inner square is k , where $0 < k < 1$. The spaces between squares are alternately shaded, as shown in the figure (which is not necessarily drawn to scale).



The area of the shaded portion of the figure is 64% of the area of the original square. What is k ?

- A $\frac{3}{5}$
- B $\frac{16}{25}$
- C $\frac{2}{3}$
- D $\frac{3}{4}$**
- E $\frac{4}{5}$

Solution:

Let the outer square have area 1. The nested squares have areas $1, k^2, k^4, \dots$, so the ring between the n th and $(n + 1)$ th squares has area $k^{2n}(1 - k^2)$.

The shaded rings are the alternate ones $n = 0, 2, 4, \dots$, with total area

$$\begin{aligned}\sum_{j=0}^{\infty} k^{4j}(1 - k^2) &= \frac{1 - k^2}{1 - k^4} \\ &= \frac{1}{1 + k^2}.\end{aligned}$$

Setting $\frac{1}{1 + k^2} = \frac{16}{25}$ gives $1 + k^2 = \frac{25}{16}$, so $k^2 = \frac{9}{16}$ and $k = \frac{3}{4}$.

Thus, the correct answer is **D**.

6. Six chairs are arranged around a round table. Two students and two teachers randomly select four of the chairs to sit in. What is the probability that the two students will sit in two adjacent chairs and the two teachers will also sit in two adjacent chairs?

A $\frac{1}{6}$

B $\frac{1}{5}$

C $\frac{2}{9}$

D $\frac{3}{13}$

E $\frac{1}{4}$

Solution:

Choosing 2 chairs for the students and 2 for the teachers gives $\binom{6}{2} \binom{4}{2} = 15 \cdot 6 = 90$ equally likely outcomes.

A round table has 6 adjacent pairs of chairs. Give the students any adjacent pair; among the remaining 4 chairs there are exactly 3 adjacent pairs for the teachers. That is $6 \cdot 3 = 18$ favorable outcomes.

The probability is $\frac{18}{90} = \frac{1}{5}$.

Thus, the correct answer is **B**.

7. In a certain alien world, the maximum running speed v of an organism is dependent on its number of toes n and number of eyes m . The relationship can be expressed as $v = kn^am^b$ centimeters per hour, where k , a , and b are integer constants. In a population where all organisms have 5 toes, $\log v = 4 + 2 \log m$; and in a population where all organisms have 25 eyes, $\log v = 4 + 4 \log n$, where the logarithms are base 10. What is $k + a + b$?

A 20

B 21

C 22

D 23

E 24

Solution:

Taking logarithms, $\log v = \log k + a \log n + b \log m$.

With $n = 5$, this reads $\log v = (\log k + a \log 5) + b \log m$, matching $4 + 2 \log m$, so $b = 2$ and $\log k + a \log 5 = 4$.

With $m = 25$, it reads $\log v = (\log k + b \log 25) + a \log n$, matching $4 + 4 \log n$, so $a = 4$ and $\log k + 2 \log 25 = 4$.

Then $\log k = 4 - \log 625 = \log \frac{10000}{625} = \log 16$, so $k = 16$. Hence $k + a + b = 16 + 4 + 2 = 22$.

Thus, the correct answer is **C**.

8. Pentagon $ABCDE$ is inscribed in a circle, and $\angle BEC = \angle CED = 30^\circ$. Let AC and BD intersect at point F , and suppose that $AB = 9$ and $AD = 24$. What is BF ?

A $\frac{57}{11}$

B $\frac{59}{11}$

C $\frac{60}{11}$

D $\frac{61}{11}$

E $\frac{63}{11}$

Solution:

The inscribed angle $\angle BEC = 30^\circ$ subtends arc $BC = 60^\circ$, so $\angle BAC$, which also subtends arc BC , equals 30° . Likewise $\angle CAD = 30^\circ$.

Thus AC bisects $\angle BAD = 60^\circ$. In $\triangle ABD$,

$$\begin{aligned} BD^2 &= 9^2 + 24^2 \\ &\quad - 2(9)(24) \cos 60^\circ \\ &= 657 - 216 = 441, \end{aligned}$$

so $BD = 21$.

Since AF (along AC) bisects $\angle BAD$, the Angle Bisector Theorem gives $\frac{BF}{FD} = \frac{AB}{AD} = \frac{9}{24} = \frac{3}{8}$. Hence $BF = \frac{3}{11} \cdot 21 = \frac{63}{11}$.

Thus, the correct answer is **E**.

9. Let w be the complex number $2 + i$, where $i = \sqrt{-1}$. What real number r has the property that r , w , and w^2 are three collinear points in the complex plane?

A $\frac{3}{4}$

B 1

C $\frac{7}{5}$

D $\frac{3}{2}$

E $\frac{5}{3}$

Solution:


Compute $w^2 = (2 + i)^2 = 3 + 4i$, so the points are $(2, 1)$ and $(3, 4)$.

The line through them has slope $\frac{4 - 1}{3 - 2} = 3$, giving $y = 3x - 5$. Setting $y = 0$ yields $x = \frac{5}{3}$.

So $r = \frac{5}{3}$.

Thus, the correct answer is **E**.

10. In the figure shown below, major arc AD and minor arc BC have the same center, O . Also, A lies between O and B , and D lies between O and C . Major arc AD , minor arc BC , and each of the two segments AB and CD have length 2π .



What is the distance from O to A ?

- A 1
- B $1 - \pi + \sqrt{1 + \pi^2}$**
- C $\frac{1}{2}\pi$
- D $\frac{1}{2}\sqrt{1 + \pi^2}$
- E 2

Solution:

Let $R_1 = OA = OD$ and $R_2 = OB = OC$, and let $\alpha = \angle AOD = \angle BOC$ (the rays coincide). The minor arc BC has length $R_2\alpha = 2\pi$, and the major arc AD is the reflex arc, so $R_1(2\pi - \alpha) = 2\pi$.

Each segment $AB = CD = R_2 - R_1 = 2\pi$.

From the first two equations, $R_2 = \frac{2\pi}{\alpha}$ and $R_1 = \frac{2\pi}{2\pi - \alpha}$. Substituting into $R_2 - R_1 = 2\pi$ and dividing by 2π gives

$$\frac{1}{\alpha} - \frac{1}{2\pi - \alpha} = 1,$$

which simplifies to $\alpha^2 - 2(1 + \pi)\alpha + 2\pi = 0$.

The larger root is greater than 2π , so it cannot be an angle of this minor arc. Thus $\alpha = (1 + \pi) - \sqrt{1 + \pi^2}$. Then

$$\begin{aligned} R_1 &= \frac{2\pi}{2\pi - \alpha} \\ &= \frac{2\pi}{\pi - 1 + \sqrt{1 + \pi^2}} \\ &= 1 - \pi + \sqrt{1 + \pi^2}, \end{aligned}$$

after rationalizing (the denominator times $\sqrt{1 + \pi^2} - (\pi - 1)$ equals 2π).

Thus, the correct answer is **B**.

11. The orthocenter of a triangle is the concurrent intersection of the three (possibly extended) altitudes. What is the sum of the coordinates of the orthocenter of the triangle whose vertices are $A(2, 31)$, $B(8, 27)$, and $C(18, 27)$?

A 5

B 17

C $10 + 4\sqrt{17} + 2\sqrt{13}$

D $\frac{113}{3}$

E 54

Solution:

Since B and C both have $y = 27$, side BC is horizontal and the altitude from A is the vertical line $x = 2$.

Side AC has slope $\frac{27 - 31}{18 - 2} = -\frac{1}{4}$, so the altitude from B has slope 4:

$$y - 27 = 4(x - 8).$$

At $x = 2$, $y = 27 + 4(2 - 8) = 3$. The orthocenter is $(2, 3)$, with coordinate sum 5.

Thus, the correct answer is **A**.

12. The harmonic mean of a collection of numbers is the reciprocal of the arithmetic mean of the reciprocals of the numbers in the collection. For example, the harmonic mean of 4, 4, and 5 is

$$\frac{1}{\frac{1}{3} \left(\frac{1}{4} + \frac{1}{4} + \frac{1}{5} \right)} = \frac{30}{7}.$$

What is the harmonic mean of all the real roots of the 4050th degree polynomial

$$\begin{aligned} & \prod_{k=1}^{2025} (kx^2 - 4x - 3) \\ &= (x^2 - 4x - 3) \\ & \quad \cdot (2x^2 - 4x - 3) \\ & \quad \cdot (3x^2 - 4x - 3) \cdots \\ & \quad (2025x^2 - 4x - 3)? \end{aligned}$$

- A $-\frac{5}{3}$
- B $-\frac{3}{2}$**
- C $-\frac{6}{5}$
- D $-\frac{5}{6}$
- E $-\frac{2}{3}$

Solution:

Each factor $kx^2 - 4x - 3$ has discriminant $16 + 12k > 0$, so it has two real roots; there are 4050 roots in all.

For the roots of $kx^2 - 4x - 3$, the sum of reciprocals is $\frac{\text{sum}}{\text{product}} = \frac{\frac{4}{k}}{-\frac{3}{k}} = -\frac{4}{3}$, independent of k .

Summing over all 2025 factors, $\sum \frac{1}{r} = 2025 \left(-\frac{4}{3} \right) = -2700$. The harmonic mean is

$$\frac{4050}{-2700} = -\frac{3}{2}.$$

Thus, the correct answer is **B**.

13. Let $C = \{1, 2, 3, \dots, 13\}$. Let N be the greatest integer such that there exists a subset of C with N elements that does not contain five consecutive integers. Suppose N integers are chosen at random from C without replacement. What is the probability that the chosen elements do not include five consecutive integers?

A $\frac{3}{130}$

B $\frac{3}{143}$

C $\frac{5}{143}$

D $\frac{1}{26}$

E $\frac{5}{78}$

Solution:

To avoid five consecutive integers, it suffices to remove two elements (for example 5 and 10), and no single removal can hit both disjoint blocks $\{1, \dots, 5\}$ and $\{9, \dots, 13\}$. Thus $N = 11$.


Choosing 11 of 13 elements is the same as removing 2, which can be done in $\binom{13}{2} = 78$ ways. The chosen set avoids five consecutive integers exactly when the two removed elements together intersect every window $\{t, t + 1, t + 2, t + 3, t + 4\}$ for $t = 1, \dots, 9$.

This forces one removed element in $\{1, \dots, 5\}$, the other in $\{9, \dots, 13\}$, and the two within 5 of each other. The valid removals are $\{4, 9\}$, $\{5, 9\}$, and $\{5, 10\}$, giving 3 of them.

The probability is $\frac{3}{78} = \frac{1}{26}$.

Thus, the correct answer is **D**.

14. Points F , G , and H are collinear with G between F and H . The ellipse with foci at G and H is internally tangent to the ellipse with foci at F and G , as shown below.



The two ellipses have the same eccentricity e , and the ratio of their areas is 2025. (Recall that the eccentricity of an ellipse is $e = \frac{c}{a}$, where c is the distance from the center to a focus, and $2a$ is the length of the major axis.) What is e ?

- A $\frac{3}{5}$
- B $\frac{16}{25}$
- C $\frac{4}{5}$
- D $\frac{22}{23}$**
- E $\frac{44}{45}$

Solution:

With the same eccentricity, $b = a\sqrt{1 - e^2}$, so the area $\pi ab \propto a^2$. The area ratio 2025 gives $\frac{a_1}{a_2} = \sqrt{2025} = 45$, where a_1, a_2 are the semi-major axes.

Both ellipses share focus G . On the large ellipse G is the right focus, so its right vertex lies $a_1 - c_1$ to the right of G . On the small ellipse G is the left focus, so its right vertex lies $a_2 + c_2$ to the right of G . Internal tangency makes these coincide:

$$a_1 - c_1 = a_2 + c_2.$$

Using $c = ea$, $a_1(1 - e) = a_2(1 + e)$, so $45(1 - e) = 1 + e$, giving $46e = 44$ and $e = \frac{22}{23}$.

Thus, the correct answer is **D**.

15. A set of numbers is called sum-free if whenever x and y are (not necessarily distinct) elements of the set, $x + y$ is not an element of the set. For example, $\{1, 4, 6\}$ and the empty set are sum-free, but $\{2, 4, 5\}$ is not. What is the greatest possible number of elements in a sum-free subset of $\{1, 2, 3, \dots, 20\}$?

- ☐ A 8
- ☐ B 9
- ☒ C 10
- ☐ D 11
- ☐ E 12

Solution:

The set $\{11, 12, \dots, 20\}$ has 10 elements and is sum-free, since any two elements sum to at least $22 > 20$.

For the upper bound, let $a_1 < a_2 < \dots < a_k$ be a sum-free subset. Each difference $a_k - a_i$ for $i < k$ cannot lie in S , because $(a_k - a_i) + a_i = a_k \in S$ would violate sum-freeness.

These $k - 1$ differences are distinct, lie in $\{1, \dots, 19\}$, and are disjoint from the k elements of S . So $k + (k - 1) \leq 20$, giving $k \leq 10$.

Thus, the correct answer is **C**.

16. Triangle $\triangle ABC$ has side lengths $AB = 80$, $BC = 45$, and $AC = 75$. The bisector of $\angle B$ and the altitude to side AB intersect at point P . What is BP ?

A 18

B 19

C 20

D 21

E 22

Solution:

By the Law of Cosines,

$$\begin{aligned}\cos B &= \frac{80^2 + 45^2 - 75^2}{2 \cdot 80 \cdot 45} \\ &= \frac{2800}{7200} = \frac{7}{18}.\end{aligned}$$

The altitude to AB is drawn from C , and its foot is at distance $BC \cos B = 45 \cdot \frac{7}{18} = 17.5$ from B along AB .

Along the bisector from B , the component parallel to AB is $BP \cos \frac{B}{2}$, which must reach the altitude's foot: $BP \cos \frac{B}{2} = 17.5$.

Since $\cos \frac{B}{2} = \sqrt{\frac{1 + \frac{7}{18}}{2}} = \sqrt{\frac{25}{36}} = \frac{5}{6}$, we get $BP = \frac{17.5}{\frac{5}{6}} = 21$.

Thus, the correct answer is **D**.

17. The polynomial $(z + i)(z + 2i)(z + 3i) + 10$ has three roots in the complex plane, where $i = \sqrt{-1}$. What is the area of the triangle formed by these roots?

- ☒ A 6
- ☐ B 8
- ☐ C 10
- ☐ D 12
- ☐ E 14

Solution:

The sum of the roots is $-6i$, so the centroid is $-2i$. Substituting $z = u - 2i$,

$$\begin{aligned}(u - i)(u)(u + i) + 10 \\&= u(u^2 + 1) + 10 \\&= u^3 + u + 10.\end{aligned}$$

Since $u = -2$ is a root, $u^3 + u + 10 = (u + 2)(u^2 - 2u + 5)$, giving roots $u = -2$ and $u = 1 \pm 2i$.

These are the points $(-2, 0)$, $(1, 2)$, $(1, -2)$. The base between $(1, 2)$ and $(1, -2)$ has length 4, at horizontal distance 3 from $(-2, 0)$, so the area is $\frac{1}{2}(4)(3) = 6$. Translation does not change the area.

Thus, the correct answer is **A**.

18. How many ordered triples (x, y, z) of distinct nonnegative integers less than or equal to 8 satisfy $xy > z$, $zx > y$, and $yz > x$?

- ☐ A 36
- ☐ B 84
- ☒ C 186
- ☐ D 336
- ☐ E 486

Solution:

If any variable is 0, say $z = 0$, then $zx = 0 > y$ is impossible. So $x, y, z \in \{1, \dots, 8\}$ are distinct positive integers.

The conditions are symmetric. For distinct values $a < b < c$, we have $ac > b$ and $bc > a$ automatically, so the only real constraint is $ab > c$. When it holds, all 6 orderings work.

For each possible smallest value a , the numbers of pairs $a < b < c \leq 8$ satisfying $ab > c$ are

a	1	2	3	4	5	6
count	0	11	10	6	3	1.

For fixed a, b , this count comes from choosing c between $b + 1$ and $\min(8, ab - 1)$. Thus there are 31 unordered triples, and all 6 orderings of each work. The answer is $6 \cdot 31 = 186$.

Thus, the correct answer is **C**.

19. Let a , b , and c be the roots of the polynomial $x^3 + kx + 1$. What is the sum

$$a^3b^2 + a^2b^3 + b^3c^2 + b^2c^3 + c^3a^2 + c^2a^3?$$

A $-k$

B $-k + 1$

C 1

D $k - 1$

E k

Solution:

By Vieta's formulas, $a + b + c = 0$, $ab + bc + ca = k$, and $abc = -1$.

Group the sum as

$$a^2b^2(a + b) + b^2c^2(b + c) + c^2a^2(c + a).$$


Since $a + b + c = 0$, we have $a + b = -c$, $b + c = -a$, $c + a = -b$.

So the sum equals

$$\begin{aligned} & -a^2b^2c - ab^2c^2 - a^2bc^2 \\ & = -abc(ab + bc + ca) \\ & = -(-1)(k) = k. \end{aligned}$$

Thus, the correct answer is **E**.

20. The base of the pentahedron shown below is a 13×8 rectangle, and its lateral faces are two isosceles triangles with base of length 8 and congruent sides of length 13, and two isosceles trapezoids with bases of lengths 7 and 13 and nonparallel sides of length 13.



What is the volume of the pentahedron?

- ☐ A 416
- ☐ B 520
- ☒ C 528
- ☐ D 676
- ☐ E 832

Solution:

The top is a ridge of length 7, centered above the base at some height h . Its endpoints sit above $(3, 4)$ and $(10, 4)$ of the 13×8 base. A slant edge to a base corner has length $\sqrt{3^2 + 4^2 + h^2} = 13$, so $h = 12$.

At height z , the horizontal cross-section is a rectangle measuring $\left(13 - \frac{z}{2}\right)$ by $\left(8 - \frac{2z}{3}\right)$. At $z = 0$ its area is 104; at $z = 6$ it is $10 \cdot 4 = 40$; at $z = 12$ the ridge has area 0.

By the prismatoid formula,

$$\begin{aligned} V &= \frac{12}{6} (104 + 4 \cdot 40 + 0) \\ &= 2(264) = 528. \end{aligned}$$

Thus, the correct answer is **C**.

21. There is a unique ordered triple (a, k, m) of nonnegative integers such that

$$\frac{4^a + 4^{a+k} + 4^{a+2k} + \dots + 4^{a+mk}}{2^a + 2^{a+k} + 2^{a+2k} + \dots + 2^{a+mk}} = 964.$$

What is $a + k + m$?

- A 8
- B 9
- C 10
- D 11
- E 12

Solution:

If $k = 0$ or $m = 0$, the original ratio is just 2^a , which cannot equal 964. Hence $k, m > 0$.

Summing the geometric series, the numerator is $4^a \frac{4^{k(m+1)} - 1}{4^k - 1}$ and the denominator is $2^a \frac{2^{k(m+1)} - 1}{2^k - 1}$. Using $4^N - 1 = (2^N - 1)(2^N + 1)$, the ratio simplifies to

$$2^a \cdot \frac{2^{k(m+1)} + 1}{2^k + 1} = 964.$$

The fraction on the left is odd over odd, so its power of 2 is 0. Since $964 = 2^2 \cdot 241$, we must have $a = 2$, and

$$2^{k(m+1)} + 1 = 241(2^k + 1).$$

Reducing this equation modulo 2^k shows that $2^k \mid 240$, so $k \leq 4$. For $k = 1, 2, 3$, the required powers of 2 would be 722, 1204, 2168, none of which is a power of 2. For $k = 4$, however, $241(2^4 + 1) = 4097 = 2^{12} + 1$, so $k(m + 1) = 12$ and $m = 2$. This also proves uniqueness.

Then $a + k + m = 2 + 4 + 2 = 8$.

Thus, the correct answer is **A**.

22. Three real numbers are chosen independently and uniformly at random between 0 and 1. What is the probability that the greatest of these three numbers is greater than 2 times each of the other two numbers? (In other words, if the chosen numbers are $a \geq b \geq c$, then $a > 2b$.)

A $\frac{1}{12}$

B $\frac{1}{9}$

C $\frac{1}{8}$

D $\frac{1}{6}$

E $\frac{1}{4}$

Solution:

Order the values as $x_1 > x_2 > x_3$; the joint density of the order statistics is 6 on this region. The event is $x_1 > 2x_2$.

Integrating x_3 from 0 to x_2 contributes a factor of x_2 . Then

$$\begin{aligned} P &= 6 \int_0^{\frac{1}{2}} x_2 \int_{2x_2}^1 dx_1 dx_2 \\ &= 6 \int_0^{\frac{1}{2}} x_2 (1 - 2x_2) dx_2. \end{aligned}$$

This equals $6 \left(\frac{1}{8} - \frac{1}{12} \right) = 6 \cdot \frac{1}{24} = \frac{1}{4}$.

Thus, the correct answer is **E**.

23. Call a positive integer fair if no digit is used more than once, it has no 0s, and no digit is adjacent to two greater digits. For example, 196, 23, and 12463 are fair, but 1546, 320, and 34321 are not fair. How many fair positive integers are there?

- A 511
- B 2584
- C 9841**
- D 17711
- E 19682

Solution:

The digits are distinct and drawn from $\{1, \dots, 9\}$, and “no digit adjacent to two greater digits” means no interior digit is smaller than both neighbors.


For a fixed set of n digits, build the arrangement by inserting digits from largest to smallest; each new (smaller) digit must go to one of the two ends, giving 2^{n-1} valid arrangements.

Summing over all nonempty digit subsets,

$$\begin{aligned} & \sum_{n=1}^9 \binom{9}{n} 2^{n-1} \\ &= \frac{1}{2} \sum_{n=1}^9 \binom{9}{n} 2^n \\ &= \frac{3^9 - 1}{2} = \frac{19682}{2} = 9841. \end{aligned}$$

Thus, the correct answer is **C**.

24. A circle of radius r is surrounded by 12 circles of radius 1, externally tangent to the central circle and sequentially tangent to each other, as shown. Then r can be written as $\sqrt{a} + \sqrt{b} + c$, where a , b , and c are integers. What is $a + b + c$?



- A 3
- B 5
- C 7**
- D 9
- E 11

Solution:

The centers of the 12 outer circles lie on a circle of radius $r + 1$, forming a regular 12-gon. Adjacent centers are 2 apart (both circles have radius 1), and the central angle between them is 30° .

Thus $2(r + 1) \sin 15^\circ = 2$, so $r + 1 = \frac{1}{\sin 15^\circ}$. Since $\sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$,

$$r + 1 = \frac{4}{\sqrt{6} - \sqrt{2}} = \sqrt{6} + \sqrt{2}.$$

Then $r = \sqrt{6} + \sqrt{2} - 1$, so $a + b + c = 6 + 2 - 1 = 7$.

Thus, the correct answer is **C**.

25. Polynomials $P(x)$ and $Q(x)$ each have degree 3 and leading coefficient 1, and their roots are all elements of $\{1, 2, 3, 4, 5\}$. The function $f(x) = \frac{P(x)}{Q(x)}$ has the property that there exist real numbers $a < b < c < d$ such that the set of all real numbers x such that $f(x) \leq 0$ consists of the closed interval $[a, b]$ together with the open interval (c, d) . How many functions $f(x)$ are possible?

- A 7
- B 9
- C 11
- D 12
- E 13**

Solution:

This problem was voided: as written, its answer is not among the choices. Here is the count.

For $\{f \leq 0\} = [a, b] \cup (c, d)$, the endpoints a, b of the closed interval must be zeros of P at which $Q \neq 0$, while c, d must be poles. The required sign pattern is therefore that of

$$g(x) = \frac{(x-a)(x-b)}{(x-c)(x-d)}.$$

The unused third factor of P must match the unused third factor of Q ; otherwise there would be an extra zero, pole, or sign change.

If the common factor is the fifth value not among a, b, c, d , it cannot make a hole inside either interval. In the ordered list of five values it may therefore occur before a , between b and c , or after d , giving 3 functions.

Alternatively, the common factor can equal c or d . For each of the $\binom{5}{4} = 5$ choices of a, b, c, d , these two polynomial pairs simplify to the same formula g and have the same domain (both omit c, d), so they define only one function. This gives 5 more functions, for a literal total of $3 + 5 = 8$.

Counting the two polynomial pairs separately in each of the last five cases gives $3 + 2 \cdot 5 = 13$, the provisional answer (E), but that does not answer the stated question about functions. Thus none of the printed choices is correct.

Problems: <https://live.poshenloh.com/past-contests/amc12/2025A>

